

## Arc Length in Polar Coordinates

If a curve is given in polar coordinates  $r = f(\theta)$ , an integral for the length of the curve can be derived using the arc length formula for a parametric curve. Regard  $\theta$  as the parameter. The parametric arc length formula becomes

$$L = \int_a^b \sqrt{\left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2} d\theta.$$

Now  $x = r \cos \theta$  and  $y = r \sin \theta$ , so

$$\frac{dx}{d\theta} = -r \sin \theta + \left(\frac{dr}{d\theta}\right) \cos \theta,$$

$$\frac{dy}{d\theta} = r \cos \theta + \left(\frac{dr}{d\theta}\right) \sin \theta,$$

Square and add, using the fact that  $(\cos \theta)^2 + (\sin \theta)^2 = 1$ :


$$\begin{aligned} \left(\frac{dx}{d\theta}\right)^2 &= r^2(\sin \theta)^2 - 2r \left(\frac{dr}{d\theta}\right) \sin \theta \cos \theta + \left(\frac{dr}{d\theta}\right)^2 (\cos \theta)^2 \\ \left(\frac{dy}{d\theta}\right)^2 &= r^2(\cos \theta)^2 + 2r \left(\frac{dr}{d\theta}\right) \cos \theta \sin \theta + \left(\frac{dr}{d\theta}\right)^2 (\sin \theta)^2 \\ \hline \left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2 &= r^2 + \left(\frac{dr}{d\theta}\right)^2 \end{aligned}$$

Hence,

$$L = \int_a^b \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta.$$

Note: As with other arc length computations, it's pretty easy to come up with polar curves which lead to integrals with non-elementary antiderivatives. In that case, the best you might be able to do is to approximate the integral using a calculator or a computer.

**Example.** Find the length of the curve  $r = \theta^2 - 1$  from  $\theta = 1$  to  $\theta = 2$ .



$$\frac{dr}{d\theta} = 2\theta.$$


$$r^2 + \left(\frac{dr}{d\theta}\right)^2 = (\theta^2 - 1)^2 + 4\theta^2 = \theta^4 - 2\theta^2 + 1 + 4\theta^2 = \theta^4 + 2\theta^2 + 1 = (\theta^2 + 1)^2.$$

$$\sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} = \theta^2 + 1.$$

The length is

$$\int_1^2 (\theta^2 + 1) d\theta = \left[\frac{1}{3}\theta^3 + \theta\right]_1^2 = \frac{10}{3} = 3.33333\dots \quad \square$$

**Example.** Find the length of the cardioid  $r = 1 + \sin \theta$  for  $\theta = 0$  to  $\theta = \frac{\pi}{2}$ .



$$\frac{dr}{d\theta} = \cos \theta.$$

$$r^2 + \left(\frac{dr}{d\theta}\right)^2 = (1 + \sin \theta)^2 + (\cos \theta)^2 = 1 + 2 \sin \theta + (\sin \theta)^2 + (\cos \theta)^2 = 2 + 2 \sin \theta.$$

$$\sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} = \sqrt{2}\sqrt{1 + \sin \theta}.$$

I'll do the antiderivative separately:

$$\int \sqrt{2}\sqrt{1 + \sin \theta} d\theta = \sqrt{2} \int \frac{\sqrt{1 + \sin \theta}\sqrt{1 - \sin \theta}}{\sqrt{1 - \sin \theta}} d\theta = \sqrt{2} \int \frac{\sqrt{1 - (\sin \theta)^2}}{\sqrt{1 - \sin \theta}} d\theta =$$

$$\sqrt{2} \int \frac{\sqrt{(\cos \theta)^2}}{\sqrt{1 - \sin \theta}} d\theta = \sqrt{2} \int \frac{\cos \theta}{\sqrt{1 - \sin \theta}} d\theta =$$


$$\left[ u = 1 - \sin \theta, \quad du = -\cos \theta d\theta, \quad d\theta = \frac{du}{-\cos \theta} \right]$$

$$\sqrt{2} \int \frac{\cos \theta}{\sqrt{u}} \cdot \frac{du}{-\cos \theta} = -\sqrt{2} \int \frac{1}{\sqrt{u}} du = -\sqrt{2} \cdot 2\sqrt{u} + c = -2\sqrt{2}\sqrt{1 - \sin \theta} + c.$$

The length is

$$\int_0^{\pi/2} \sqrt{2}\sqrt{1 + \sin \theta} d\theta = \left[-2\sqrt{2}\sqrt{1 - \sin \theta}\right]_0^{\pi/2} = 2\sqrt{2} = 2.82842\dots \quad \square$$

**Example.** Find the length of the polar curve  $r = \left(\sin \frac{\theta}{4}\right)^4$  for  $\theta = 0$  to  $\theta = \pi$ .



$$\frac{dr}{d\theta} = \left(\sin \frac{\theta}{4}\right)^3 \cos \frac{\theta}{4}.$$

$$r^2 + \left(\frac{dr}{d\theta}\right)^2 = \left(\sin \frac{\theta}{4}\right)^8 + \left(\sin \frac{\theta}{4}\right)^6 \left(\cos \frac{\theta}{4}\right)^2 = \left(\sin \frac{\theta}{4}\right)^6 \left[\left(\sin \frac{\theta}{4}\right)^2 + \left(\cos \frac{\theta}{4}\right)^2\right] = \left(\sin \frac{\theta}{4}\right)^6.$$

$$\sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} = \left(\sin \frac{\theta}{4}\right)^3.$$

The length is

$$\int_0^\pi \left(\sin \frac{\theta}{4}\right)^3 d\theta.$$

I'll do the antiderivative separately:

$$\int \left(\sin \frac{\theta}{4}\right)^3 d\theta = \int \left(\sin \frac{\theta}{4}\right)^2 \sin \frac{\theta}{4} d\theta = \int \left(1 - \left(\cos \frac{\theta}{4}\right)^2\right) \sin \frac{\theta}{4} d\theta =$$

$$\left[ u = \cos \frac{\theta}{4}, \quad du = -\frac{1}{4} \sin \frac{\theta}{4} d\theta, \quad d\theta = -4 \frac{du}{\sin \frac{\theta}{4}} \right]$$

$$-4 \int (1 - u^2) \left(\sin \frac{\theta}{4}\right) \cdot \frac{du}{\sin \frac{\theta}{4}} = -4 \int (1 - u^2) du = -4 \left(u - \frac{1}{3}u^3\right) + c = -4 \cos \frac{\theta}{4} + \frac{4}{3} \left(\cos \frac{\theta}{4}\right)^3 + c.$$

So

$$\int_0^\pi \left(\sin \frac{\theta}{4}\right)^3 d\theta = \left[-4 \cos \frac{\theta}{4} + \frac{4}{3} \left(\cos \frac{\theta}{4}\right)^3\right]_0^\pi = \frac{8}{3} - \frac{5\sqrt{2}}{3} = 0.30964\dots \quad \square$$