

# Introduction to Fourier Series

## MATH 467 *Partial Differential Equations*

J. Robert Buchanan

Department of Mathematics

Fall 2018

# Objectives

In this lesson we will learn:

- ▶ the formal process for finding a Fourier series representation of a function,
- ▶ the orthogonality of the trigonometric functions,
- ▶ the Euler-Fourier formulas for finding Fourier series coefficients,
- ▶ properties of periodic functions,
- ▶ how to periodically extend a function,
- ▶ the properties of even and odd periodic extensions of functions, and
- ▶ practice finding the Fourier series representations of functions.

# Informal Definition of a Fourier Series

The **Fourier series** expansion of a function  $f(x)$  is a representation of  $f(x)$  on an interval  $[-L, L]$  as the sum of sine and cosine functions of the form

$$\frac{a_0}{2} + \sum_{n=1}^{\infty} \left( a_n \cos \frac{n\pi x}{L} + b_n \sin \frac{n\pi x}{L} \right)$$

where  $a_n$  and  $b_n$  are constants.

## Issues Raised by Fourier Series

- ▶ What functions  $f(x)$  can be written as a Fourier series?
- ▶ If  $f(x)$  can be represented as a Fourier Series, what are the constants  $a_n$  and  $b_n$ ?
- ▶ Will the Fourier series converge?
- ▶ Provided the Fourier series converges, does it converge to  $f(x)$  at all points in the interval  $[-L, L]$ ?
- ▶ Can Fourier series be differentiated and integrated?

# Inner Product

## Definition

If  $u(x)$  and  $v(x)$  are integrable on  $[a, b]$ , the **inner product** of  $u$  and  $v$  on  $[a, b]$ , denoted as  $\langle u, v \rangle$ , is defined as

$$\langle u, v \rangle = \int_a^b u(x)v(x) \, dx.$$

## Definition

The functions  $u$  and  $v$  are said to be **orthogonal** on  $[a, b]$  if

$$\langle u, v \rangle = \int_a^b u(x)v(x) \, dx = 0.$$

A set  $S$  of integrable functions on  $[a, b]$  is said to be a **mutually orthogonal set** if each pair of distinct functions in the set is orthogonal.

# Trigonometric System

Let  $S$  be the infinite set of functions

$$\left\{ 1, \cos \frac{\pi x}{L}, \sin \frac{\pi x}{L}, \cos \frac{2\pi x}{L}, \sin \frac{2\pi x}{L}, \dots, \cos \frac{n\pi x}{L}, \sin \frac{n\pi x}{L}, \dots \right\}.$$

$S$  is a mutually orthogonal set on  $[-L, L]$ .

## Product-to-Sum Formulas

$$\cos \alpha \cos \beta = \frac{1}{2} (\cos(\alpha + \beta) + \cos(\alpha - \beta))$$

$$\cos \alpha \sin \beta = \frac{1}{2} (\sin(\alpha + \beta) - \sin(\alpha - \beta))$$

$$\sin \alpha \sin \beta = \frac{1}{2} (\cos(\alpha - \beta) - \cos(\alpha + \beta))$$

## Justification of Orthogonality

$$\begin{aligned}
 & \int_{-L}^L \cos \frac{m\pi x}{L} \cos \frac{n\pi x}{L} dx \\
 &= \frac{1}{2} \int_{-L}^L \left[ \cos \frac{(m+n)\pi x}{L} + \cos \frac{(m-n)\pi x}{L} \right] dx \\
 &= \begin{cases} \frac{L}{2\pi} \left[ \frac{1}{m+n} \sin \frac{(m+n)\pi x}{L} + \frac{1}{m-n} \sin \frac{(m-n)\pi x}{L} \right]_{-L}^L & \text{if } m \neq n, \\ \frac{1}{2} \left[ \frac{L}{2m\pi} \sin \frac{2m\pi x}{L} + x \right]_{-L}^L & \text{if } m = n \end{cases} \\
 &= \begin{cases} 0 & \text{if } m \neq n, \\ L & \text{if } m = n. \end{cases}
 \end{aligned}$$

The orthogonality of  $\sin(m\pi x/L)$ ,  $\sin(n\pi x/L)$ , and  $\cos(k\pi x/L)$  is handled similarly.

# Euler-Fourier Formulas

Assuming  $f(x)$  defined on  $[-L, L]$  can be represented as a Fourier series we write

$$f(x) \sim \frac{a_0}{2} + \sum_{n=1}^{\infty} \left( a_n \cos \frac{n\pi x}{L} + b_n \sin \frac{n\pi x}{L} \right),$$

where

$$a_0 = \frac{1}{L} \int_{-L}^L f(x) dx$$

$$a_n = \frac{1}{L} \int_{-L}^L f(x) \cos \frac{n\pi x}{L} dx$$

$$b_n = \frac{1}{L} \int_{-L}^L f(x) \sin \frac{n\pi x}{L} dx$$

for  $n = 1, 2, \dots$

## Justification (1 of 2)

Assuming  $f(x)$  equals its Fourier representation on  $[-L, L]$  and that the infinite series can be integrated term-by-term, multiply both sides of the equation

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left( a_n \cos \frac{n\pi x}{L} + b_n \sin \frac{n\pi x}{L} \right),$$

by  $\sin(m\pi x/L)$  and integrate over  $[-L, L]$ .

$$\begin{aligned} & \int_{-L}^L f(x) \sin \frac{m\pi x}{L} dx \\ &= \int_{-L}^L \left[ \frac{a_0}{2} + \sum_{n=1}^{\infty} \left( a_n \cos \frac{n\pi x}{L} + b_n \sin \frac{n\pi x}{L} \right) \right] \sin \frac{m\pi x}{L} dx \\ &= \frac{a_0}{2} \int_{-L}^L \sin \frac{m\pi x}{L} dx + \sum_{n=1}^{\infty} a_n \int_{-L}^L \cos \frac{n\pi x}{L} \sin \frac{m\pi x}{L} dx \\ &\quad + \sum_{n=1}^{\infty} b_n \int_{-L}^L \sin \frac{n\pi x}{L} \sin \frac{m\pi x}{L} dx = b_m L \end{aligned}$$

## Justification (2 of 2)

Multiplying both sides of the earlier equation by  $\cos(m\pi x/L)$  and integrating over  $[-L, L]$  yields  $a_m$  for  $m \in \mathbb{N}$ .

Integrating both sides of

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left( a_n \cos \frac{n\pi x}{L} + b_n \sin \frac{n\pi x}{L} \right),$$

over  $[-L, L]$  produces

$$\begin{aligned} \int_{-L}^L f(x) dx &= \int_{-L}^L \frac{a_0}{2} dx \\ &\quad + \sum_{n=1}^{\infty} \left( a_n \int_{-L}^L \cos \frac{n\pi x}{L} dx + b_n \int_{-L}^L \sin \frac{n\pi x}{L} dx \right) \\ &= a_0 L. \end{aligned}$$

## Remarks

- ▶ In general the symbol  $\sim$  is used in place of  $=$  since we do not yet know whether the infinite series converges, or if it does converge, that it converges to  $f(x)$ .
- ▶ The only assumption placed on  $f(x)$  is that it be integrable on  $[-L, L]$ . It does not even need to be defined at all points in  $[-L, L]$ .
- ▶ If the infinite series converges, it does so to a  $2L$ -periodic function, which can be thought of as the  $2L$ -periodic extension of  $f(x)$ .

# Periodic Functions

## Definition

A function  $f(x)$  is said to be **periodic** if there exists a constant  $T > 0$  such that, for any  $x$  in the domain of  $f$ ,  $x + T$  is in its domain and  $f(x + T) = f(x)$  holds for all such  $x$ . In this case,  $T$  is called a **period** of  $f(x)$  and, often  $f(x)$  is said to be  $T$ -**periodic** or **periodic with period  $T$** .

# Properties of Periodic Functions

- ▶ Any constant function is periodic and any  $T > 0$  is a period.
- ▶ If  $T$  is a period of function  $f(x)$ , so is  $kT$  for any  $k \in \mathbb{N}$ .
- ▶ If  $f(x)$  and  $g(x)$  are periodic with common period  $T$ , then for any constant  $c$ ,  $cf(x)$ ,  $f(x) \pm g(x)$ ,  $f(x) \cdot g(x)$ , and  $f(x)/g(x)$  are all periodic with period  $T$  on their respective domains.
- ▶ If  $f(x)$  is periodic with period  $T$ , then so is  $f'(x)$  on its domain.
- ▶ If  $f(x)$  is  $T$ -periodic, integrable and  $\int_0^T f(x) dx = 0$ , then

$$\int_0^x f(t) dt \text{ is } T\text{-periodic.}$$

- ▶ If  $f(x)$  is an integrable, periodic function with period  $T$  defined on  $(-\infty, \infty)$ , then for any  $a \in \mathbb{R}$ ,

$$\int_a^{a+T} f(x) dx = \int_0^T f(x) dx.$$

# Periodic Extensions

Suppose  $f(x)$  is defined on  $[-L, L]$  where  $L > 0$ . A periodic function  $F(x)$  can be defined on  $(-\infty, \infty)$  in the following way:


- ▶ If  $x \in (-L, L]$ , then  $F(x) = f(x)$ .
- ▶ If  $x \notin (-L, L]$  and  $k$  is an integer such that  $x + k(2L) \in (-L, L]$ , then  $F(x) = f(x + k(2L))$ .

## Remarks:

- ▶  $F(x)$  is periodic with period  $2L$ .
- ▶ If no confusion results,  $f(x)$  is used to denote its own periodic extension.
- ▶  $F(x)$  as defined not a “true” extension of  $f(x)$  unless  $f(-L) = f(L)$ .


## Example (1 of 2)

Function  $f(x) = x^2$  is continuous on  $[-1, 1]$ . Sketch its 2-periodic extension.



## Example (2 of 2)

Function  $f(x) = e^x$  is continuous on  $[-1, 1]$ . Sketch its 2-periodic extension.



# Find the Fourier Coefficients

Consider the piecewise-defined function

$$f(x) = \begin{cases} x & \text{if } -1 \leq x < 0, \\ 0 & \text{if } 0 \leq x < 1. \end{cases}$$

1. Write down the Fourier series of  $f(x)$ .
2. Sketch the 2-periodic extension of  $f(x)$ .


## Coefficients

$$\begin{aligned}a_0 &= \frac{1}{1} \int_{-1}^1 f(x) dx = \int_{-1}^0 x dx = -\frac{1}{2} \\a_n &= \frac{1}{1} \int_{-1}^1 f(x) \cos \frac{n\pi x}{1} dx = \int_{-1}^0 x \cos(n\pi x) dx \\&= \frac{1 - (-1)^n}{n^2 \pi^2} = \begin{cases} 2/(n\pi)^2 & \text{if } n \text{ is odd,} \\ 0 & \text{if } n \text{ is even.} \end{cases} \\b_n &= \frac{1}{1} \int_{-1}^1 f(x) \sin \frac{n\pi x}{1} dx = \int_{-1}^0 x \sin(n\pi x) dx \\&= \frac{(-1)^{n+1}}{n\pi}\end{aligned}$$


# Fourier Representation

$$f(x) \sim -\frac{1}{4} + \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n\pi} \sin(n\pi x) + \sum_{n=1}^{\infty} \frac{2}{(2n-1)^2\pi^2} \cos((2n-1)\pi x)$$

## 2-Periodic Extension



## Fourier Series (truncated to 10 terms)



## Find the Fourier Coefficients

Consider the function  $f(x) = x^2$ .


1. Write down the Fourier series of  $f(x)$  valid for  $[-\pi, \pi]$ .
2. Sketch the  $2\pi$ -periodic extension of  $f(x)$ .

# Coefficients


$$\begin{aligned}a_0 &= \frac{1}{\pi} \int_{-\pi}^{\pi} x^2 dx = \frac{2}{3} \pi^2 \\a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} x^2 \cos(nx) dx = \frac{4(-1)^n}{n^2} \\b_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} x^2 \sin(nx) dx = 0\end{aligned}$$

$$f(x) \sim \frac{\pi^2}{3} + \sum_{n=1}^{\infty} \frac{4(-1)^n}{n^2} \cos(nx)$$

## $2\pi$ -Periodic Extension



## Fourier Series (truncated to 10 terms)



## Find the Fourier Coefficients

Consider the function

$$f(x) = \begin{cases} 0 & \text{if } -\pi \leq x \leq 0, \\ \sin x & \text{if } 0 < x < \pi. \end{cases}$$

1. Write down the Fourier series of  $f(x)$  valid for  $[-\pi, \pi]$ .
2. Sketch the  $2\pi$ -periodic extension of  $f(x)$ .

# Coefficients

$$a_0 = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) dx = \frac{1}{\pi} \int_0^{\pi} \sin x dx = \frac{2}{\pi}$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos(nx) dx = \frac{1}{\pi} \int_0^{\pi} \sin x \cos(nx) dx$$


$$= \begin{cases} -2/(\pi(n^2 - 1)) & \text{if } n \text{ is even,} \\ 0 & \text{if } n \text{ is odd.} \end{cases}$$

$$b_1 = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin x dx = \frac{1}{\pi} \int_0^{\pi} \sin^2 x dx = \frac{1}{2}$$


$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin(nx) dx = \frac{1}{\pi} \int_0^{\pi} \sin x \sin(nx) dx = 0$$

$$f(x) \sim \frac{1}{\pi} + \frac{1}{2} \sin x - \sum_{n=1}^{\infty} \frac{2}{\pi(4n^2 - 1)} \cos(2nx)$$

## $2\pi$ -Periodic Extension



## Fourier Series (truncated to 10 terms)



## Find the Fourier Coefficients

Find the Fourier series representation of  $g(x) = |\sin x|$  on  $[-\pi, \pi]$ .

# Solution


- ▶ Note that

$$|\sin x| = -\sin x + \begin{cases} 0 & \text{if } -\pi \leq x \leq 0, \\ 2\sin x & \text{if } 0 \leq x \leq \pi. \end{cases}$$

- ▶ The Fourier series for  $\sin x$  is merely  $\sin x$ .
- ▶ The Fourier series for the piecewise-defined function was found in the previous example.

$$f(x) \sim \frac{2}{\pi} - \sum_{n=1}^{\infty} \frac{4}{\pi(4n^2 - 1)} \cos(2nx)$$

## Fourier Series (truncated to 10 terms)



## Even, Odd, Periodic Extensions

**Comment:** the spatial domain of many of the PDEs we study (e.g., the heat equation and wave equation) is the interval  $[0, L]$ , not  $[-L, L]$ . If an initial condition is specified on  $[0, L]$  we may extend it to  $[-L, L]$  (and thence to  $(-\infty, \infty)$ ) in any way that it remains integrable. Options include:

### Even Extension

$$f_e(x) = \begin{cases} f(-x) & \text{if } -L \leq x < 0, \\ f(x) & \text{if } 0 \leq x \leq L. \end{cases}$$

### Odd Extension


$$f_o(x) = \begin{cases} -f(-x) & \text{if } -L \leq x < 0, \\ f(x) & \text{if } 0 \leq x \leq L. \end{cases}$$

## Example

Consider the function  $f(x) = \cos x$  on  $[0, \pi/2]$ .

1. Sketch the odd  $\pi$ -periodic extension of  $f(x)$ .
2. Find the Fourier series representation for the odd  $\pi$ -periodic extension of  $f(x)$ .

Graph of  $f_o(x)$



## Fourier Series Coefficients

$$a_0 = \frac{2}{\pi} \int_{-\pi/2}^{\pi/2} f_0(x) dx = 0$$

$$a_n = \frac{2}{\pi} \int_{-\pi/2}^{\pi/2} f_0(x) \cos \frac{n\pi x}{\pi/2} dx = 0$$

$$b_n = \frac{2}{\pi} \int_{-\pi/2}^{\pi/2} f_0(x) \sin \frac{n\pi x}{\pi/2} dx$$


$$= -\frac{2}{\pi} \int_{-\pi/2}^0 \cos(-x) \sin \frac{n\pi x}{\pi/2} dx + \frac{2}{\pi} \int_0^{\pi/2} \cos(x) \sin \frac{n\pi x}{\pi/2} dx$$

$$= \frac{4}{\pi} \int_0^{\pi/2} \cos(x) \sin(2nx) dx = \frac{8n}{(4n^2 - 1)\pi}$$

Since only the  $b_n$  coefficients are nonzero, this is called a **Fourier sine series**.

# Fourier Series Representation

$$f_o(x) \sim \sum_{n=1}^{\infty} \frac{8n}{(4n^2 - 1)\pi} \sin(2nx)$$




## Example

Consider the function  $f(x) = e^x$  on  $[0, 1]$ .

1. Sketch the even 2-periodic extension of  $f(x)$ .
2. Find the Fourier series representation for the even 2-periodic extension of  $f(x)$ .

Graph of  $f_e(x)$




## Fourier Series Coefficients

$$\begin{aligned}a_0 &= \int_{-1}^1 f_e(x) dx \\&= 2 \int_0^1 e^x dx = 2(e - 1) \\a_n &= \int_{-1}^1 f_e(x) \cos(n\pi x) dx \\&= 2 \int_0^1 e^x \cos(n\pi x) dx = \frac{2((-1)^n e - 1)}{n^2 \pi^2 + 1} \\b_n &= \int_{-1}^1 f_e(x) \sin(n\pi x) dx = 0\end{aligned}$$

Since only the  $a_n$  coefficients are nonzero, this is called a **Fourier cosine series**.

# Fourier Series Representation

$$f_e(x) \sim e - 1 + \sum_{n=1}^{\infty} \frac{2((-1)^n e - 1)}{n^2 \pi^2 + 1} \cos(n\pi x)$$



## Remark

Any function  $f(x)$  defined on  $(-\infty, \infty)$  can be written as the sum of an even function and an odd function. In fact,

$$f(x) = \frac{f(x) + f(-x)}{2} + \frac{f(x) - f(-x)}{2}$$

where  $(f(x) + f(-x))/2$  is even (sometimes called the **even part** of  $f$ ) and  $(f(x) - f(-x))/2$  is odd (likewise called the **odd part** of  $f$ ).

# Homework

- ▶ Read Sections 3.1–3.5
- ▶ Exercises: 1–9