

<div data-bbox="220 1379 304 1912" data-label="Section-Header"> <h1>Introduction to Fourier Series</h1> <h2>MATH 467 Partial Differential Equations</h2> </div> <div data-bbox="379 1520 545 1769" data-label="Text"> <p>J Robert Buchanan Department of Mathematics Fall 2022</p> </div>	<div data-bbox="25 902 65 1093" data-label="Section-Header"> <h3>Objectives</h3> </div> <div data-bbox="189 694 217 1034" data-label="Text"> <p>In this lesson we will learn:</p> </div> <div data-bbox="233 147 595 1014" data-label="List-Group"> <ul style="list-style-type: none"> ▶ the formal process for finding a Fourier series representation of a function, ▶ the orthogonality of the trigonometric functions, ▶ the Euler-Fourier formulas for finding Fourier series coefficients, ▶ properties of periodic functions, ▶ how to periodically extend a function, ▶ the properties of even and odd periodic extensions of functions, and ▶ practice finding the Fourier series representations of functions. </div>
<div data-bbox="821 1460 861 2145" data-label="Section-Header"> <h3>Informal Definition of a Fourier Series</h3> </div> <div data-bbox="1061 1202 1158 2092" data-label="Text"> <p>The Fourier series expansion of a function $f(x)$ is a representation of $f(x)$ on an interval $[-L, L]$ as the sum of sine and cosine functions of the form</p> </div> <div data-bbox="1160 1424 1243 1872" data-label="Equation-Block"> $\frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi x}{L} + b_n \sin \frac{n\pi x}{L} \right)$ </div> <div data-bbox="1264 1693 1291 2092" data-label="Text"> <p>where a_n and b_n are constants.</p> </div>	<div data-bbox="821 506 861 1093" data-label="Section-Header"> <h3>Issues Raised by Fourier Series</h3> </div> <div data-bbox="1045 174 1319 1014" data-label="List-Group"> <ul style="list-style-type: none"> ▶ What functions $f(x)$ can be written as a Fourier series? ▶ If $f(x)$ can be represented as a Fourier Series, what are the constants a_n and b_n? ▶ Will the Fourier series converge? ▶ Provided the Fourier series converges, does it converge to $f(x)$ at all points in the interval $[-L, L]$? ▶ Can Fourier series be differentiated and integrated? </div>

Inner Product

Definition

If $u(x)$ and $v(x)$ are integrable on $[a, b]$, the **inner product** of u and v on $[a, b]$, denoted as $\langle u, v \rangle$, is defined as

$$\langle u, v \rangle = \int_a^b u(x)v(x) \, dx.$$

Definition

The functions u and v are said to be **orthogonal** on $[a, b]$ if

$$\langle u, v \rangle = \int_a^b u(x)v(x) \, dx = 0.$$

A set S of integrable functions on $[a, b]$ is said to be a **mutually orthogonal set** if each pair of distinct functions in the set is orthogonal.

Trigonometric System

Let S be the infinite set of functions

$$\left\{ 1, \cos \frac{\pi x}{L}, \sin \frac{\pi x}{L}, \cos \frac{2\pi x}{L}, \sin \frac{2\pi x}{L}, \dots, \cos \frac{n\pi x}{L}, \sin \frac{n\pi x}{L}, \dots \right\}.$$

S is a mutually orthogonal set on $[-L, L]$.

Product-to-Sum Formulas

$$\begin{aligned} \cos \alpha \cos \beta &= \frac{1}{2} (\cos(\alpha + \beta) + \cos(\alpha - \beta)) \\ \cos \alpha \sin \beta &= \frac{1}{2} (\sin(\alpha + \beta) - \sin(\alpha - \beta)) \\ \sin \alpha \sin \beta &= \frac{1}{2} (\cos(\alpha - \beta) - \cos(\alpha + \beta)) \end{aligned}$$

Justification of Orthogonality

$$\begin{aligned} &\int_{-L}^L \cos \frac{m\pi x}{L} \cos \frac{n\pi x}{L} \, dx \\ &= \frac{1}{2} \int_{-L}^L \left[\cos \frac{(m+n)\pi x}{L} + \cos \frac{(m-n)\pi x}{L} \right] \, dx \\ &= \begin{cases} \frac{L}{2\pi} \left[\frac{1}{m+n} \sin \frac{(m+n)\pi x}{L} + \frac{1}{m-n} \sin \frac{(m-n)\pi x}{L} \right]_{-L}^L & \text{if } m \neq n, \\ \frac{1}{2} \left[\frac{L}{2m\pi} \sin \frac{2m\pi x}{L} + x \right]_{-L}^L & \text{if } m = n \end{cases} \\ &= \begin{cases} 0 & \text{if } m \neq n, \\ L & \text{if } m = n. \end{cases} \end{aligned}$$

The orthogonality of $\sin(m\pi x/L)$, $\sin(n\pi x/L)$, and $\cos(k\pi x/L)$ is handled similarly.

Euler-Fourier Formulas

Assuming $f(x)$ defined on $[-L, L]$ can be represented as a Fourier series we write

$$f(x) \sim \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi x}{L} + b_n \sin \frac{n\pi x}{L} \right),$$

where

$$\begin{aligned} a_0 &= \frac{1}{L} \int_{-L}^L f(x) dx \\ a_n &= \frac{1}{L} \int_{-L}^L f(x) \cos \frac{n\pi x}{L} dx \\ b_n &= \frac{1}{L} \int_{-L}^L f(x) \sin \frac{n\pi x}{L} dx \end{aligned}$$

for $n = 1, 2, \dots$

Justification (2 of 2)

Multiplying both sides of the earlier equation by $\cos(m\pi x/L)$ and integrating over $[-L, L]$ yields a_m for $m \in \mathbb{N}$.

Integrating both sides of

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi x}{L} + b_n \sin \frac{n\pi x}{L} \right),$$

over $[-L, L]$ produces

$$\begin{aligned} \int_{-L}^L f(x) dx &= \int_{-L}^L \frac{a_0}{2} dx + \sum_{n=1}^{\infty} \left(a_n \int_{-L}^L \cos \frac{n\pi x}{L} dx + b_n \int_{-L}^L \sin \frac{n\pi x}{L} dx \right) \\ &= a_0 L. \end{aligned}$$

Justification (1 of 2)

Assuming $f(x)$ equals its Fourier representation on $[-L, L]$ and that the infinite series can be integrated term-by-term, multiply both sides of the equation

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi x}{L} + b_n \sin \frac{n\pi x}{L} \right),$$

by $\sin(m\pi x/L)$ and integrate over $[-L, L]$.

$$\begin{aligned} \int_{-L}^L f(x) \sin \frac{m\pi x}{L} dx &= \int_{-L}^L \left[\frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos \frac{n\pi x}{L} + b_n \sin \frac{n\pi x}{L} \right) \right] \sin \frac{m\pi x}{L} dx \\ &= \frac{a_0}{2} \int_{-L}^L \sin \frac{m\pi x}{L} dx + \sum_{n=1}^{\infty} a_n \int_{-L}^L \cos \frac{n\pi x}{L} \sin \frac{m\pi x}{L} dx \\ &\quad + \sum_{n=1}^{\infty} b_n \int_{-L}^L \sin \frac{n\pi x}{L} \sin \frac{m\pi x}{L} dx = b_m L \end{aligned}$$

Remarks

- ▶ In general the symbol \sim is used in place of $=$ since we do not yet know whether the infinite series converges, or if it does converge, that it converges to $f(x)$.
- ▶ The only assumption placed on $f(x)$ is that it be integrable on $[-L, L]$. It does not even need to be defined at all points in $[-L, L]$.
- ▶ If the infinite series converges, it does so to a $2L$ -periodic function, which can be thought of as the $2L$ -periodic extension of $f(x)$.

Periodic Functions

Definition

A function $f(x)$ is said to be **periodic** if there exists a constant $T > 0$ such that, for any x in the domain of f , $x + T$ is in its domain and $f(x + T) = f(x)$ holds for all such x . In this case, T is called a **period** of $f(x)$ and, often $f(x)$ is said to be T -**periodic** or **periodic with period T** .

Properties of Periodic Functions

- ▶ Any constant function is periodic and any $T > 0$ is a period.
- ▶ If T is a period of function $f(x)$, so is kT for any $k \in \mathbb{N}$.
- ▶ If $f(x)$ and $g(x)$ are periodic with common period T , then for any constant c , $cf(x)$, $f(x) \pm g(x)$, $f(x) \cdot g(x)$, and $f(x)/g(x)$ are all periodic with period T on their respective domains.
- ▶ If $f(x)$ is periodic with period T , then so is $f'(x)$ on its domain.
- ▶ If $f(x)$ is T -periodic, integrable and $\int_0^T f(x) dx = 0$, then $\int_0^x f(t) dt$ is T -periodic.
- ▶ If $f(x)$ is an integrable, periodic function with period T defined on $(-\infty, \infty)$, then for any $a \in \mathbb{R}$,

$$\int_a^{a+T} f(x) dx = \int_0^T f(x) dx.$$

Periodic Extensions

Suppose $f(x)$ is defined on $[-L, L]$ where $L > 0$. A periodic function $F(x)$ can be defined on $(-\infty, \infty)$ in the following way:


- ▶ If $x \in (-L, L]$, then $F(x) = f(x)$.
- ▶ If $x \notin (-L, L]$ and k is an integer such that $x + k(2L) \in (-L, L]$, then $F(x) = f(x + k(2L))$.

Remarks:

- ▶ $F(x)$ is periodic with period $2L$.
- ▶ If no confusion results, $f(x)$ is used to denote its own periodic extension.
- ▶ $F(x)$ as defined not a "true" extension of $f(x)$ unless $f(-L) = f(L)$.


Example (1 of 2)

Function $f(x) = x^2$ is continuous on $[-1, 1]$. Sketch its 2-periodic extension.



Example (2 of 2)

Function $f(x) = e^x$ is continuous on $[-1, 1]$. Sketch its 2-periodic extension.



Find the Fourier Coefficients

Consider the piecewise-defined function

$$f(x) = \begin{cases} x & \text{if } -1 \leq x < 0, \\ 0 & \text{if } 0 \leq x < 1. \end{cases}$$

1. Write down the Fourier series of $f(x)$.
2. Sketch the 2-periodic extension of $f(x)$.

Coefficients

$$a_0 = \frac{1}{1} \int_{-1}^1 f(x) dx = \int_{-1}^0 x dx = -\frac{1}{2}$$

$$a_n = \frac{1}{1} \int_{-1}^1 f(x) \cos \frac{n\pi x}{1} dx = \int_{-1}^0 x \cos(n\pi x) dx$$

$$= \frac{1 - (-1)^n}{n^2 \pi^2} = \begin{cases} 2/(n\pi)^2 & \text{if } n \text{ is odd,} \\ 0 & \text{if } n \text{ is even.} \end{cases}$$


$$b_n = \frac{1}{1} \int_{-1}^1 f(x) \sin \frac{n\pi x}{1} dx = \int_{-1}^0 x \sin(n\pi x) dx$$

$$= \frac{(-1)^{n+1}}{n\pi}$$


Fourier Representation

$$f(x) \sim -\frac{1}{4} + \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n\pi} \sin(n\pi x) + \sum_{n=1}^{\infty} \frac{2}{(2n-1)^2 \pi^2} \cos((2n-1)\pi x)$$

2-Periodic Extension



Fourier Series (truncated to 10 terms)




Find the Fourier Coefficients

- Consider the function $f(x) = x^2$.
1. Write down the Fourier series of $f(x)$ valid for $[-\pi, \pi]$.
 2. Sketch the 2π -periodic extension of $f(x)$.


Coefficients

$$a_0 = \frac{1}{\pi} \int_{-\pi}^{\pi} x^2 dx = \frac{2}{3} \pi^2$$
$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} x^2 \cos(nx) dx = \frac{4(-1)^n}{n^2}$$
$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} x^2 \sin(nx) dx = 0$$
$$f(x) \sim \frac{\pi^2}{3} + \sum_{n=1}^{\infty} \frac{4(-1)^n}{n^2} \cos(nx)$$

2 π -Periodic Extension



Fourier Series (truncated to 10 terms)



Find the Fourier Coefficients

Consider the function

$$f(x) = \begin{cases} 0 & \text{if } -\pi \leq x \leq 0, \\ \sin x & \text{if } 0 < x < \pi. \end{cases}$$



1. Write down the Fourier series of $f(x)$ valid for $[-\pi, \pi]$.
2. Sketch the 2π -periodic extension of $f(x)$.

Coefficients

$$\begin{aligned} a_0 &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) dx = \frac{1}{\pi} \int_0^{\pi} \sin x dx = \frac{2}{\pi} \\ a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos(nx) dx = \frac{1}{\pi} \int_0^{\pi} \sin x \cos(nx) dx \\ &= \begin{cases} -2/(\pi(n^2 - 1)) & \text{if } n \text{ is even,} \\ 0 & \text{if } n \text{ is odd.} \end{cases} \\ b_1 &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin x dx = \frac{1}{\pi} \int_0^{\pi} \sin^2 x dx = \frac{1}{2} \\ b_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin(nx) dx = \frac{1}{\pi} \int_0^{\pi} \sin x \sin(nx) dx = 0 \end{aligned}$$

$$f(x) \sim \frac{1}{\pi} + \frac{1}{2} \sin x - \sum_{n=1}^{\infty} \frac{2}{\pi(4n^2 - 1)} \cos(2nx)$$

2π-Periodic Extension



Fourier Series (truncated to 10 terms)

Find the Fourier Coefficients

Find the Fourier series representation of $g(x) = |\sin x|$ on $[-\pi, \pi]$.

Solution


► Note that

$$|\sin x| = -\sin x + \begin{cases} 0 & \text{if } -\pi \leq x \leq 0, \\ 2 \sin x & \text{if } 0 \leq x \leq \pi. \end{cases}$$

- The Fourier series for $\sin x$ is merely $\sin x$.
- The Fourier series for the piecewise-defined function was found in the previous example.

$$f(x) \sim \frac{2}{\pi} - \sum_{n=1}^{\infty} \frac{4}{\pi(4n^2 - 1)} \cos(2nx)$$

Fourier Series (truncated to 10 terms)



Even, Odd, Periodic Extensions

Comment: the spatial domain of many of the PDEs we study (e.g., the heat equation and wave equation) is the interval $[0, L]$, not $[-L, L]$. If an initial condition is specified on $[0, L]$ we may extend it to $[-L, L]$ (and thence to $(-\infty, \infty)$) in any way that it remains integrable. Options include:

Even Extension

$$f_e(x) = \begin{cases} f(-x) & \text{if } -L \leq x < 0, \\ f(x) & \text{if } 0 \leq x \leq L. \end{cases}$$

Odd Extension


$$f_o(x) = \begin{cases} -f(-x) & \text{if } -L \leq x < 0, \\ f(x) & \text{if } 0 \leq x \leq L. \end{cases}$$

Example

Consider the function $f(x) = \cos x$ on $[0, \pi/2]$.

1. Sketch the odd π -periodic extension of $f(x)$.
2. Find the Fourier series representation for the odd π -periodic extension of $f(x)$.

Graph of $f_o(x)$



Fourier Series Coefficients

$$a_0 = \frac{2}{\pi} \int_{-\pi/2}^{\pi/2} f_o(x) dx = 0$$

$$a_n = \frac{2}{\pi} \int_{-\pi/2}^{\pi/2} f_o(x) \cos \frac{n\pi x}{\pi/2} dx = 0$$


$$b_n = \frac{2}{\pi} \int_{-\pi/2}^{\pi/2} f_o(x) \sin \frac{n\pi x}{\pi/2} dx$$

$$\begin{aligned} &= -\frac{2}{\pi} \int_{-\pi/2}^0 \cos(-x) \sin \frac{n\pi x}{\pi/2} dx + \frac{2}{\pi} \int_0^{\pi/2} \cos(x) \sin \frac{n\pi x}{\pi/2} dx \\ &= \frac{4}{\pi} \int_0^{\pi/2} \cos(x) \sin(2nx) dx = \frac{8n}{(4n^2 - 1)\pi} \end{aligned}$$

Since only the b_n coefficients are nonzero, this is called a **Fourier sine series**.

Fourier Series Representation

$$f_o(x) \sim \sum_{n=1}^{\infty} \frac{8n}{(4n^2 - 1)\pi} \sin(2nx)$$




Example

Consider the function $f(x) = e^x$ on $[0, 1]$.

1. Sketch the even 2-periodic extension of $f(x)$.
2. Find the Fourier series representation for the even 2-periodic extension of $f(x)$.

Graph of $f_e(x)$




Fourier Series Coefficients

$$\begin{aligned}
 a_0 &= \int_{-1}^1 f_e(x) dx \\
 &= 2 \int_0^1 e^x dx = 2(e - 1) \\
 a_n &= \int_{-1}^1 f_e(x) \cos(n\pi x) dx \\
 &= 2 \int_0^1 e^x \cos(n\pi x) dx = \frac{2((-1)^n e - 1)}{n^2 \pi^2 + 1} \\
 b_n &= \int_{-1}^1 f_e(x) \sin(n\pi x) dx = 0
 \end{aligned}$$

Since only the a_n coefficients are nonzero, this is called a **Fourier cosine series**.

Fourier Series Representation

$$f_e(x) \sim e - 1 + \sum_{n=1}^{\infty} \frac{2((-1)^n e - 1)}{n^2 \pi^2 + 1} \cos(n\pi x)$$



Remark

Any function $f(x)$ defined on $(-\infty, \infty)$ can be written as the sum of an even function and an odd function. In fact,

$$f(x) = \frac{f(x) + f(-x)}{2} + \frac{f(x) - f(-x)}{2}$$

where $(f(x) + f(-x))/2$ is even (sometimes called the **even part** of f) and $(f(x) - f(-x))/2$ is odd (likewise called the **odd part** of f).

Homework

- ▶ Read Sections 3.1–3.5
- ▶ Exercises: 1–9