

<div data-bbox="220 1323 260 1968" data-label="Section-Header"> <h1>Neumann Problems on Rectangles</h1> </div> <div data-bbox="272 1391 303 1901" data-label="Text"> <p>MATH 467 <i>Partial Differential Equations</i></p> </div> <div data-bbox="379 1520 406 1771" data-label="Text"> <p>J Robert Buchanan</p> </div> <div data-bbox="451 1520 472 1771" data-label="Text"> <p>Department of Mathematics</p> </div> <div data-bbox="518 1583 545 1702" data-label="Text"> <p>Fall 2022</p> </div>	<div data-bbox="25 902 65 1095" data-label="Section-Header"> <h2>Objectives</h2> </div> <div data-bbox="266 692 293 1037" data-label="Text"> <p>In this lesson we will learn:</p> </div> <div data-bbox="311 199 485 1014" data-label="List-Group"> <ul style="list-style-type: none"> <li>▶ to solve Laplace's equation on two-dimensional domains with Neumann boundary conditions,</li> <li>▶ to compare the solutions on domains with Dirichlet boundary conditions to solution on domains with Neumann boundary conditions.</li> </ul> </div>
<div data-bbox="821 1688 861 2148" data-label="Section-Header"> <h2>Boundary Value Problem</h2> </div> <div data-bbox="973 1319 1070 2092" data-label="Text"> <p>Consider Laplace's equation on the rectangle <math>\Omega = \{(x, y) \mid 0 &lt; x &lt; a, 0 &lt; y &lt; b\}</math> with Neumann boundary conditions:</p> </div> <div data-bbox="1107 1411 1268 1881" data-label="Equation-Block"> <math display="block">\begin{aligned} \Delta u &amp;= 0 \text{ for } (x, y) \in \Omega \\ u_y(x, 0) &amp;= u_y(x, b) = 0 \text{ for } 0 &lt; x &lt; a \\ u_x(0, y) &amp;= 0 \text{ for } 0 &lt; y &lt; b \\ u_x(a, y) &amp;= f(y) \text{ for } 0 &lt; y &lt; b. \end{aligned}</math> </div> <div data-bbox="1324 1207 1422 2092" data-label="Text"> <p><b>Physical Interpretation:</b> the steady-state heat distribution in <math>\Omega</math> when the rectangular region is insulated along its bottom, top, and left edges and there is a flow of heat on the right edge.</p> </div>	<div data-bbox="821 768 861 1095" data-label="Section-Header"> <h2>Product Solutions</h2> </div> <div data-bbox="916 176 979 1037" data-label="Text"> <p>Assume <math>u(x, y) = X(x)Y(y)</math> solves Laplace's equation. Separation of variables induces the following ODEs for <math>X(x)</math> and <math>Y(y)</math>.</p> </div> <div data-bbox="1013 315 1088 869" data-label="Equation-Block"> <math display="block">\begin{aligned} X''(x) - \sigma X(x) &amp;= 0 \text{ with } X'(0) = 0 \\ Y''(y) + \sigma Y(y) &amp;= 0 \text{ with } Y'(0) = 0 = Y'(b), \end{aligned}</math> </div> <div data-bbox="1121 754 1149 1037" data-label="Text"> <p>where <math>\sigma</math> is a constant.</p> </div> <div data-bbox="1177 320 1208 1037" data-label="Text"> <p>Taking the second BVP, the only nontrivial solutions are:</p> </div> <div data-bbox="1238 488 1374 694" data-label="Equation-Block"> <math display="block">\begin{aligned} Y_n(y) &amp;= \cos \frac{n\pi y}{b} \\ \sigma_n &amp;= \frac{n^2\pi^2}{b^2} \end{aligned}</math> </div> <div data-bbox="1404 813 1431 1037" data-label="Text"> <p>for <math>n = 0, 1, 2, \dots</math></p> </div> <div data-bbox="1453 416 1513 1037" data-label="Text"> <p>This implies <math>X_n(x) = \cosh \frac{n\pi x}{b}</math> for <math>n = 0, 1, 2, \dots</math></p> </div>

## Series Solution

The product solutions:

$$u_n(x, y) = \cosh \frac{n\pi x}{b} \cos \frac{n\pi y}{b} \text{ for } n = 0, 1, 2, \dots$$

solve Laplace's equation and satisfy the three homogeneous boundary conditions.

By the Principle of Superposition,

$$u(x, y) = b_0 + \sum_{n=1}^{\infty} b_n \cosh \frac{n\pi x}{b} \cos \frac{n\pi y}{b}.$$

The coefficients  $b_n$  can be determined from the remaining nonhomogeneous boundary condition.

## Determining the Coefficients

Differentiate the formal series solution and let  $x = a$ .

$$u_x(a, y) = \sum_{n=1}^{\infty} b_n \left( \frac{n\pi}{b} \right) \sinh \frac{n\pi a}{b} \cos \frac{n\pi y}{b} = f(y)$$

**Remarks:**

- ▶ The coefficient  $b_0$  was lost during the differentiation.
- ▶ The infinite series can be regarded as a cosine series for  $f(y)$  if the integral of  $f$  over  $[0, b]$  vanishes, i.e., if

$$\int_0^b f(y) dy = 0.$$

## Further Remarks

- ▶ If  $\int_0^b f(y) dy \neq 0$  then a solution to the BVP does not exist.
- ▶ Consider the physics:
  - ▶ If the definite integral vanishes then there is no net flux of heat across the boundary at  $x = a$  and hence a steady-state (time independent) heat distribution can evolve.
  - ▶ If the definite integral does not vanish, then there is a net flux of heat in or out of  $\Omega$  and no time independent temperature distribution can exist.
- ▶ Even if the definite integral vanishes, the solution can be determined only up to the addition of an arbitrary constant. Thus Laplace's equation on a rectangle with Neumann boundary conditions on all four edges has no unique solution.
- ▶ This type of boundary value problem is ill-posed.

Assuming  $\int_0^b f(y) dy = 0$

$$b_n = \frac{2}{n\pi \sinh \frac{n\pi a}{b}} \int_0^b f(y) \cos \frac{n\pi y}{b} dy,$$

for  $n \in \mathbb{N}$ .

$$u(x, y) = b_0 + \sum_{n=1}^{\infty} b_n \cosh \frac{n\pi x}{b} \cos \frac{n\pi y}{b}$$

where  $b_0$  is an arbitrary constant.

## Example

Find a solution to the Neumann boundary value problem on the unit square:

$$\Delta u = 0 \text{ for } 0 < x < 1 \text{ and } 0 < y < 1$$

$$u_y(x, 0) = u_y(x, 1) = 0 \text{ for } 0 < x < 1$$

$$u_x(0, y) = 0 \text{ for } 0 < y < 1$$

$$u_x(1, y) = y - 1/2 \text{ for } 0 < y < 1.$$

## Solution (1 of 2)

Check:  $\int_0^1 \left( y - \frac{1}{2} \right) dy = \left[ \frac{y^2}{2} - \frac{y}{2} \right]_{y=0}^{y=1} = 0.$


Using the Euler-Fourier coefficient formula:

$$\begin{aligned} b_n &= \frac{2}{n\pi \sinh(n\pi)} \int_0^1 \left( y - \frac{1}{2} \right) \cos(n\pi y) dy \\ &= \frac{-2}{n^2\pi^2 \sinh(n\pi)} \int_0^1 \sin(n\pi y) dy \\ &= \frac{2((-1)^n - 1)}{n^3\pi^3 \sinh(n\pi)}. \end{aligned}$$

$$u(x, y) = b_0 - \frac{4}{\pi^3} \sum_{n=1}^{\infty} \frac{\cosh((2n-1)\pi x) \cos((2n-1)\pi y)}{(2n-1)^3 \sinh((2n-1)\pi)}$$

where  $b_0$  is an arbitrary constant.

## Solution (2 of 2)



## General Case

Consider Laplace's equation on a rectangle with Neumann BCs on all four edges.

$$\begin{aligned} u_{xx} + u_{yy} &= 0 \text{ for } (x, y) \in R \\ u_x(0, y) &= g_1(y) \text{ for } 0 < y < b \\ u_x(a, y) &= g_2(y) \text{ for } 0 < y < b \\ u_y(x, 0) &= f_1(x) \text{ for } 0 < x < a \\ u_y(x, b) &= f_2(x) \text{ for } 0 < x < a \end{aligned}$$

This BVP can be decomposed into four sub-problems with homogeneous boundary conditions on three edges.

## Sub-Problems

$$\begin{aligned}
 \Delta u_1 &= 0 \text{ for } (x, y) \in R \\
 (u_1)_x(0, y) &= g_1(y) \text{ for } y \in (0, b) \\
 (u_1)_x(a, y) &= 0 \text{ for } y \in (0, b) \\
 (u_1)_y(x, 0) &= 0 \text{ for } x \in (0, a) \\
 (u_1)_y(x, b) &= 0 \text{ for } x \in (0, a) \\
 \\
 \Delta u_3 &= 0 \text{ for } (x, y) \in R \\
 (u_3)_x(0, y) &= 0 \text{ for } y \in (0, b) \\
 (u_3)_x(a, y) &= 0 \text{ for } y \in (0, b) \\
 (u_3)_y(x, 0) &= f_1(y) \text{ for } x \in (0, a) \\
 (u_3)_y(x, b) &= 0 \text{ for } x \in (0, a)
 \end{aligned}$$

$$\begin{aligned}
 \Delta u_2 &= 0 \text{ for } (x, y) \in R \\
 (u_2)_x(0, y) &= 0 \text{ for } y \in (0, b) \\
 (u_2)_x(a, y) &= g_2(y) \text{ for } y \in (0, b) \\
 (u_2)_y(x, 0) &= 0 \text{ for } x \in (0, a) \\
 (u_2)_y(x, b) &= 0 \text{ for } x \in (0, a) \\
 \\
 \Delta u_4 &= 0 \text{ for } (x, y) \in R \\
 (u_4)_x(0, y) &= 0 \text{ for } y \in (0, b) \\
 (u_4)_x(a, y) &= 0 \text{ for } y \in (0, b) \\
 (u_4)_y(x, 0) &= 0 \text{ for } x \in (0, a) \\
 (u_4)_y(x, b) &= f_2(y) \text{ for } x \in (0, a)
 \end{aligned}$$

$$\begin{aligned}
 u_1(x, y) &= a_0 + \sum_{n=1}^{\infty} a_n \cosh \frac{n\pi(a-x)}{b} \cos \frac{n\pi y}{b} \\
 u_2(x, y) &= b_0 + \sum_{n=1}^{\infty} b_n \cosh \frac{n\pi x}{b} \cos \frac{n\pi y}{b} \\
 u_3(x, y) &= c_0 + \sum_{n=1}^{\infty} c_n \cos \frac{n\pi x}{a} \cosh \frac{n\pi(b-y)}{a} \\
 u_4(x, y) &= d_0 + \sum_{n=1}^{\infty} d_n \cos \frac{n\pi x}{a} \cosh \frac{n\pi y}{a}
 \end{aligned}$$

## Solutions to the Sub-Problems

## Series Coefficients

Provided  $\int_0^b g_1(y) dy = \int_0^b g_2(y) dy = 0$  and  $\int_0^a f_1(x) dx = \int_0^a f_2(x) dx = 0$ , then

$$\begin{aligned}
 a_n &= \frac{-2}{n\pi \sinh \frac{n\pi a}{b}} \int_0^b g_1(y) \cos \frac{n\pi y}{b} dy \\
 b_n &= \frac{2}{n\pi \sinh \frac{n\pi a}{b}} \int_0^b g_2(y) \cos \frac{n\pi y}{b} dy \\
 c_n &= \frac{-2}{n\pi \sinh \frac{n\pi b}{a}} \int_0^a f_1(x) \cos \frac{n\pi x}{a} dx \\
 d_n &= \frac{2}{n\pi \sinh \frac{n\pi b}{a}} \int_0^a f_2(x) \cos \frac{n\pi x}{a} dx.
 \end{aligned}$$

The solution to the original BVP is then

$$u(x, y) = u_1(x, y) + u_2(x, y) + u_3(x, y) + u_4(x, y).$$

## Neumann Problems on Disks

Consider Laplace's equation on the disk of radius  $a > 0$ :

$$\Delta u = 0 \text{ for } x^2 + y^2 < a^2$$

$$\frac{\partial u}{\partial \mathbf{n}}(x, y) = \phi(x, y) \text{ for } x^2 + y^2 = a^2.$$

$\partial u / \partial \mathbf{n}$  denotes the derivative in the direction of the unit outward normal vector to the boundary.

Convert to polar coordinates.

$$\begin{aligned}
 v_{rr} + \frac{1}{r} v_r + \frac{1}{r^2} v_{\theta\theta} &= 0 \text{ for } 0 < r < a \text{ and } -\infty < \theta < \infty \\
 \frac{\partial v}{\partial r}(a, \theta) &= \phi(a \cos \theta, a \sin \theta) = f(\theta) \text{ for } -\infty < \theta < \infty.
 \end{aligned}$$

## Series Solution

The formal series solution can be written as

$$v(r, \theta) = d_0 + \sum_{n=1}^{\infty} r^n [c_n \cos(n\theta) + d_n \sin(n\theta)],$$

with coefficients  $d_0$ ,  $c_n$ , and  $d_n$  chosen such that

$$v_r(a, \theta) = \sum_{n=1}^{\infty} na^{n-1} [c_n \cos(n\theta) + d_n \sin(n\theta)] = f(\theta).$$

A necessary condition for the solution to exist is

$$\int_{-\pi}^{\pi} f(\theta) d\theta = \int_{-\pi}^{\pi} \phi(a \cos \theta, a \sin \theta) d\theta = 0.$$

## Series Coefficients

$$c_n = \frac{a^{1-n}}{n\pi} \int_{-\pi}^{\pi} f(\theta) \cos(n\theta) d\theta$$

$$d_n = \frac{a^{1-n}}{n\pi} \int_{-\pi}^{\pi} f(\theta) \sin(n\theta) d\theta.$$

**Remark:** Coefficient  $d_0$  can be chosen arbitrarily and thus the solution to Laplace's equation on a disk with Neumann boundary conditions is not unique.

## Example

Find a bounded solution to Laplace's equation on  $\Omega = \{(r, \theta) \mid 0 \leq r < 1\}$  that satisfies the Neumann boundary condition,

$$u_r(1, \theta) = f(\theta) = \theta,$$

## Solution (1 of )

The solution can be written as

$$u(r, \theta) = d_0 + \sum_{n=1}^{\infty} r^n [c_n \cos(n\theta) + d_n \sin(n\theta)].$$

The boundary condition implies

$$u_r(1, \theta) = \sum_{n=1}^{\infty} n c_n \cos(n\theta) + n d_n \sin(n\theta) = \theta.$$

Applying the Euler-Fourier formula:


$$n c_n = \frac{1}{\pi} \int_{-\pi}^{\pi} \theta \cos(n\theta) d\theta = 0$$

$$n d_n = \frac{1}{\pi} \int_{-\pi}^{\pi} \theta \sin(n\theta) d\theta$$

$$= \left[ \frac{-\theta}{n\pi} \cos(n\theta) \right]_{\theta=-\pi}^{\theta=\pi} + \frac{1}{n\pi} \int_{-\pi}^{\pi} \cos(n\theta) d\theta$$

$$d_n = \frac{-2(-1)^n}{n^2}.$$

$$u(r, \theta) = d_0 - 2 \sum_{n=1}^{\infty} \frac{(-1)^n r^n}{n^2} \sin(n\theta).$$



- ▶ Read Sections 6.5 and 6.6
- ▶ Exercises: 20–23