Neumann Problems on Rectangles MATH 467 Partial Differential Equations

J Robert Buchanan

Department of Mathematics

Fall 2022

▲□▶ ▲□▶ ▲□▶ ▲□▶ ▲□ ● のへぐ

Objectives

In this lesson we will learn:

- to solve Laplace's equation on two-dimensional domains with Neumann boundary conditions,
- to compare the solutions on domains with Dirichlet boundary conditions to solution on domains with Neumann boundary conditions.

▲□▶▲□▶▲□▶▲□▶ □ のQ@

Boundary Value Problem

Consider Laplace's equation on the rectangle $\Omega = \{(x, y) | 0 < x < a, 0 < y < b\}$ with Neumann boundary conditions:

Physical Interpretation: the steady-state heat distribution in Ω when the rectangular region is insulated along its bottom, top, and left edges and there is a flow of heat on the right edge.

(ロ) (同) (三) (三) (三) (○) (○)

Product Solutions

Assume u(x, y) = X(x)Y(y) solves Laplace's equation. Separation of variables induces the following ODEs for X(x) and Y(y).

$$X''(x) - \sigma X(x) = 0 \text{ with } X'(0) = 0$$

$$Y''(y) + \sigma Y(y) = 0 \text{ with } Y'(0) = 0 = Y'(b),$$

where σ is a constant.

Taking the second BVP, the only nontrivial solutions are:

$$Y_n(y) = \cos \frac{n\pi y}{b}$$
 $\sigma_n = \frac{n^2 \pi^2}{b^2}$

(日) (日) (日) (日) (日) (日) (日)

for *n* = 0, 1, 2, . . .

This implies $X_n(x) = \cosh \frac{n\pi x}{b}$ for n = 0, 1, 2, ...

Series Solution

The product solutions:

$$u_n(x,y) = \cosh \frac{n\pi x}{b} \cos \frac{n\pi y}{b}$$
 for $n = 0, 1, 2, \dots$

solve Laplace's equation and satisfy the three homogeneous boundary conditions.

By the Principle of Superposition,

$$u(x,y) = b_0 + \sum_{n=1}^{\infty} b_n \cosh \frac{n\pi x}{b} \cos \frac{n\pi y}{b}.$$

(ロ) (同) (三) (三) (三) (○) (○)

The coefficients b_n can be determined from the remaining nonhomogeneous boundary condition.

Determining the Coefficients

Differentiate the formal series solution and let x = a.

$$u_x(a, y) = \sum_{n=1}^{\infty} b_n\left(\frac{n\pi}{b}\right) \sinh \frac{n\pi a}{b} \cos \frac{n\pi y}{b} = f(y)$$

Remarks:

- The coefficient b_0 was lost during the differentiation.
- The infinite series can be regarded as a cosine series for f(y) if the integral of f over [0, b] vanishes, *i.e.*, if

$$\int_0^b f(y)\,dy=0.$$

(日) (日) (日) (日) (日) (日) (日)

Further Remarks

• If $\int_0^b f(y) dy \neq 0$ then a solution to the BVP does not exist.

- Consider the physics:
 - If the definite integral vanishes then there is no net flux of heat across the boundary at x = a and hence a steady-state (time independent) heat distribution can evolve.
 - If the definite integral does not vanish, then there is a net flux of heat in or out of Ω and no time independent temperature distribution can exist.
- Even if the definite integral vanishes, the solution can be determined only up to the addition of an arbitrary constant. Thus Laplace's equation on a rectangle with Neumann boundary conditions on all four edges has no unique solution.
- This type of boundary value problem is ill-posed.

Assuming
$$\int_0^b f(y) \, dy = 0$$

$$b_n = \frac{2}{n\pi\sinh\frac{n\pi a}{b}}\int_0^b f(y)\cos\frac{n\pi y}{b}\,dy,$$

for $n \in \mathbb{N}$.

$$u(x,y) = b_0 + \sum_{n=1}^{\infty} b_n \cosh \frac{n\pi x}{b} \cos \frac{n\pi y}{b}$$

where b_0 is an arbitrary constant.

Example

Find a solution to the Neumann boundary value problem on the unit square:

$$\Delta u = 0 \text{ for } 0 < x < 1 \text{ and } 0 < y < 1$$

$$u_y(x,0) = u_y(x,1) = 0 \text{ for } 0 < x < 1$$

$$u_x(0,y) = 0 \text{ for } 0 < y < 1$$

$$u_x(1,y) = y - 1/2 \text{ for } 0 < y < 1.$$

Solution (1 of 2)

Check:
$$\int_0^1 \left(y - \frac{1}{2} \right) \, dy = \left[\frac{y^2}{2} - \frac{y}{2} \right]_{y=0}^{y=1} = 0.$$

Using the Euler-Fourier coefficient formula:

$$b_n = \frac{2}{n\pi \sinh(n\pi)} \int_0^1 \left(y - \frac{1}{2}\right) \cos(n\pi y) \, dy$$

= $\frac{-2}{n^2 \pi^2 \sinh(n\pi)} \int_0^1 \sin(n\pi y) \, dy$
= $\frac{2((-1)^n - 1)}{n^3 \pi^3 \sinh(n\pi)}.$

$$u(x,y) = b_0 - \frac{4}{\pi^3} \sum_{n=1}^{\infty} \frac{\cosh((2n-1)\pi x) \cos((2n-1)\pi y)}{(2n-1)^3 \sinh((2n-1)\pi)}$$

▲□▶ ▲□▶ ▲□▶ ▲□▶ ▲□ ● のへぐ

where b_0 is an arbitrary constant.

Solution (2 of 2)



◆□▶ ◆□▶ ◆三▶ ◆三▶ ・三 の々で

General Case

Consider Laplace's equation on a rectangle with Neumann BCs on all four edges.

$$u_{xx} + u_{yy} = 0 \text{ for } (x, y) \in R$$

$$u_x(0, y) = g_1(y) \text{ for } 0 < y < b$$

$$u_x(a, y) = g_2(y) \text{ for } 0 < y < b$$

$$u_y(x, 0) = f_1(x) \text{ for } 0 < x < a$$

$$u_y(x, b) = f_2(x) \text{ for } 0 < x < a$$

▲□▶ ▲□▶ ▲ 三▶ ▲ 三▶ - 三 - のへぐ

This BVP can be decomposed into four sub-problems with homogeneous boundary conditions on three edges.

Sub-Problems

$$\begin{array}{rcl} \triangle u_3 &=& 0 \text{ for } (x,y) \in R \\ (u_3)_x(0,y) &=& 0 \text{ for } y \in (0,b) \\ (u_3)_x(a,y) &=& 0 \text{ for } y \in (0,b) \\ (u_3)_y(x,0) &=& f_1(y) \text{ for } x \in (0,a) \\ (u_3)_y(x,b) &=& 0 \text{ for } x \in (0,a) \end{array}$$

▲□▶ ▲□▶ ▲目▶ ▲目▶ 三日 - 釣A@

Solutions to the Sub-Problems

$$u_1(x, y) = a_0 + \sum_{n=1}^{\infty} a_n \cosh \frac{n\pi(a-x)}{b} \cos \frac{n\pi y}{b}$$
$$u_2(x, y) = b_0 + \sum_{n=1}^{\infty} b_n \cosh \frac{n\pi x}{b} \cos \frac{n\pi y}{b}$$
$$u_3(x, y) = c_0 + \sum_{n=1}^{\infty} c_n \cos \frac{n\pi x}{a} \cosh \frac{n\pi(b-y)}{a}$$
$$u_4(x, y) = d_0 + \sum_{n=1}^{\infty} d_n \cos \frac{n\pi x}{a} \cosh \frac{n\pi y}{a}$$

◆□ > ◆□ > ◆ 三 > ◆ 三 > ● ○ ○ ○ ○

Series Coefficients

Provided
$$\int_0^b g_1(y) \, dy = \int_0^b g_2(y) \, dy = 0$$
 and $\int_0^a f_1(x) \, dx = \int_0^a f_2(x) \, dx = 0$, then

$$a_n = \frac{-2}{n\pi\sinh\frac{n\pi a}{b}} \int_0^b g_1(y)\cos\frac{n\pi y}{b} \, dy$$
$$b_n = \frac{2}{n\pi\sinh\frac{n\pi a}{b}} \int_0^b g_2(y)\cos\frac{n\pi y}{b} \, dy$$
$$c_n = \frac{-2}{n\pi\sinh\frac{n\pi b}{a}} \int_0^a f_1(x)\cos\frac{n\pi x}{a} \, dx$$
$$d_n = \frac{2}{n\pi\sinh\frac{n\pi b}{a}} \int_0^a f_2(x)\cos\frac{n\pi x}{a} \, dx.$$

The solution to the original BVP is then

$$u(x,y) = u_1(x,y) + u_2(x,y) + u_3(x,y) + u_4(x,y).$$

◆□ > ◆□ > ◆臣 > ◆臣 > ─臣 ─のへで

Neumann Problems on Disks

Consider Laplace's equation on the disk of radius a > 0:

▲□▶ ▲□▶ ▲ 三▶ ▲ 三▶ - 三 - のへぐ

 $\partial u/\partial \mathbf{n}$ denotes the derivative in the direction of the unit outward normal vector to the boundary.

Neumann Problems on Disks

Consider Laplace's equation on the disk of radius a > 0:

 $\partial u/\partial \mathbf{n}$ denotes the derivative in the direction of the unit outward normal vector to the boundary.

Convert to polar coordinates.

$$v_{rr} + rac{1}{r}v_r + rac{1}{r^2}v_{ heta heta} = 0 ext{ for } 0 < r < a ext{ and } -\infty < heta < \infty$$

 $rac{\partial v}{\partial r}(a, heta) = \phi(a\cos heta, a\sin heta) = f(heta) ext{ for } -\infty < heta < \infty.$

< □ > < 同 > < 三 > < 三 > < 三 > < ○ < ○ </p>

Series Solution

The formal series solution can be written as

$$v(r,\theta) = d_0 + \sum_{n=1}^{\infty} r^n [c_n \cos(n\theta) + d_n \sin(n\theta)],$$

with coefficients d_0 , c_n , and d_n chosen such that

$$v_r(a,\theta) = \sum_{n=1}^{\infty} n a^{n-1} [c_n \cos(n\theta) + d_n \sin(n\theta)] = f(\theta).$$

A necessary condition for the solution to exist is

$$\int_{-\pi}^{\pi} f(\theta) \, d\theta = \int_{-\pi}^{\pi} \phi(a\cos\theta, a\sin\theta) \, d\theta = 0.$$

◆□ ▶ ◆□ ▶ ◆三 ▶ ◆□ ▶ ◆□ ●

Series Coefficients

$$c_n = \frac{a^{1-n}}{n\pi} \int_{-\pi}^{\pi} f(\theta) \cos(n\theta) \, d\theta$$
$$d_n = \frac{a^{1-n}}{n\pi} \int_{-\pi}^{\pi} f(\theta) \sin(n\theta) \, d\theta.$$

Remark: Coefficient d_0 can be chosen arbitrarily and thus the solution to Laplace's equation on a disk with Neumann boundary conditions is not unique.

▲□▶ ▲□▶ ▲ 三▶ ▲ 三▶ - 三 - のへぐ

Example

Find a bounded solution to Laplace's equation on $\Omega = \{(r, \theta) | 0 \le r < 1\}$ that satisfies the Neumann boundary condition,

$$u_r(1,\theta)=f(\theta)=\theta,$$

▲□▶ ▲□▶ ▲□▶ ▲□▶ ▲□ ● のへぐ

Solution (1 of)

The solution can be written as

$$u(r,\theta) = d_0 + \sum_{n=1}^{\infty} r^n [c_n \cos(n\theta) + d_n \sin(n\theta)].$$

The boundary condition implies

$$u_r(1,\theta) = \sum_{n=1}^{\infty} n c_n \cos(n\theta) + n d_n \sin(n\theta)] = \theta.$$

Applying the Euler-Fourier formula:

$$n c_n = \frac{1}{\pi} \int_{-\pi}^{\pi} \theta \cos(n\theta) d\theta = 0$$

$$n d_n = \frac{1}{\pi} \int_{-\pi}^{\pi} \theta \sin(n\theta) d\theta$$

$$= \left[\frac{-\theta}{n\pi} \cos(n\theta)\right]_{\theta=-\pi}^{\theta=\pi} + \frac{1}{n\pi} \int_{-\pi}^{\pi} \cos(n\theta) d\theta$$

$$d_n = \frac{-2(-1)^n}{n^2}.$$

▲□▶ ▲□▶ ▲□▶ ▲□▶ ▲□ ● のへぐ

Solution (2 of)



Homework

Read Sections 6.5 and 6.6

Exercises: 20–23