

# Maximum Principle and Uniqueness of Solutions

MATH 467 *Partial Differential Equations*

J Robert Buchanan

Department of Mathematics

Fall 2022

# Objectives

In this lesson we will explore:

- ▶ the Maximum Principle for solutions to the heat equation and its justification,
- ▶ the dependence of solutions to the heat equation on the initial and boundary conditions, and
- ▶ the uniqueness of solutions to the heat equation and its justification.

# The Maximum Principle

## Theorem

*Consider the initial boundary value problem*


$$\begin{aligned}u_t &= k u_{xx}, & 0 < x < L, & \quad t > 0 \\u(0, t) &= a(t) \quad \text{and} \quad u(L, t) = b(t), & \quad t > 0 \\u(x, 0) &= f(x), & 0 \leq x \leq L\end{aligned}$$

*where the diffusion constant  $k > 0$ , and the functions  $a(t)$ ,  $b(t)$ , and  $f(x)$  are  $C^2$  (twice continuously differentiable) on their respective intervals. Let  $T > 0$  be a fixed time and let*

$$A = \max_{0 \leq t \leq T} \{a(t)\}, \quad B = \max_{0 \leq t \leq T} \{b(t)\}, \quad \text{and} \quad F = \max_{0 \leq x \leq L} \{f(x)\}.$$

*If  $M = \max\{A, B, F\}$  and if  $u(x, t)$  is any  $C^2$  solution of the initial boundary value problem, then  $u(x, t) \leq M$  for all  $0 \leq x \leq L$  and  $0 \leq t \leq T$ .*

# Illustration



## Example

Consider the IBVP:

$$\begin{aligned}u_t &= 9u_{xx} \text{ for } 0 < x < 3 \text{ and } t > 0 \\u(0, t) &= 0 = u(3, t) \\u(x, 0) &= 6 \sin \frac{\pi x}{3} + 2 \sin \pi x\end{aligned}$$

Find an upper bound for the solution.

## Solution (1 of 3)

- ▶ According to the Maximum Principle, the maximum occurs either where  $x = 0$ ,  $x = 3$ , or  $t = 0$ .
- ▶ Either the upper bound is  $u(0, t) = u(3, t) = 0$  or the maximum occurs where

$$u(x, 0) = f(x) = 6 \sin \frac{\pi x}{3} + 2 \sin \pi x.$$

## Solution (1 of 3)

- ▶ According to the Maximum Principle, the maximum occurs either where  $x = 0$ ,  $x = 3$ , or  $t = 0$ .
- ▶ Either the upper bound is  $u(0, t) = u(3, t) = 0$  or the maximum occurs where


$$u(x, 0) = f(x) = 6 \sin \frac{\pi x}{3} + 2 \sin \pi x.$$

$$\begin{aligned} f'(x) &= 2\pi \left( \cos \frac{\pi x}{3} + \cos \pi x \right) \\ &= 2\pi \left( \cos \left( \frac{2\pi x}{3} - \frac{\pi x}{3} \right) + \cos \left( \frac{2\pi x}{3} + \frac{\pi x}{3} \right) \right) \\ &= 4\pi \cos \frac{2\pi x}{3} \cos \frac{\pi x}{3} \end{aligned}$$

## Solution (2 of 3)

$$f'(x) = 4\pi \cos \frac{2\pi x}{3} \cos \frac{\pi x}{3} = 0$$


has critical numbers  $x = 3/4$ ,  $x = 9/4$  (both maxima) and  $x = 3/2$  (minimum) in the interval  $[0, 3]$ .





## Solution (3 of 3)

$$u(x, t) \leq f(3/4) = f(9/4) = 4\sqrt{2}$$



# Minimum Principle

## Corollary

Consider the initial boundary value problem

$$\begin{aligned}u_t &= ku_{xx}, & 0 < x < L, & \quad t > 0 \\u(0, t) &= a(t) \quad \text{and} \quad u(L, t) = b(t), & \quad t > 0 \\u(x, 0) &= f(x), & 0 \leq x \leq L\end{aligned}$$

where the diffusion constant  $k > 0$ , and the functions  $a(t)$ ,  $b(t)$ , and  $f(x)$  are  $C^2$  on their respective intervals. Let  $T > 0$  be a fixed time and let

$$\alpha = \min_{0 \leq t \leq T} \{a(t)\}, \quad \beta = \min_{0 \leq t \leq T} \{b(t)\}, \quad \text{and} \quad \gamma = \min_{0 \leq x \leq L} \{f(x)\}.$$

If  $\mu = \min\{\alpha, \beta, \gamma\}$  and if  $u(x, t)$  is any  $C^2$  solution of the initial boundary value problem, then  $u(x, t) \geq \mu$  for all  $0 \leq x \leq L$  and  $0 \leq t \leq T$ .

# Continuous Dependence on BC and IC

## Theorem

Consider the two initial boundary value problems

$$\begin{array}{ll} u_t = ku_{xx} & v_t = kv_{xx} \\ u(0, t) = a_1(t) & v(0, t) = a_2(t) \\ u(L, t) = b_1(t) & v(L, t) = b_2(t) \\ u(x, 0) = f_1(x) & v(x, 0) = f_2(x) \end{array}$$

defined for  $0 \leq x \leq L$  and  $t \geq 0$ . Let  $T > 0$  and suppose there exists  $\epsilon \geq 0$  such that

$$\begin{array}{l} |f_1(x) - f_2(x)| \leq \epsilon \text{ for } 0 \leq x \leq L, \\ |a_1(t) - a_2(t)| \leq \epsilon \text{ for } 0 \leq t \leq T, \text{ and} \\ |b_1(t) - b_2(t)| \leq \epsilon \text{ for } 0 \leq t \leq T. \end{array}$$

If  $u(x, t)$  and  $v(x, t)$  are  $C^2$  solutions respectively to the two initial boundary value problems, then for all  $0 \leq x \leq L$  and  $0 \leq t \leq T$ ,

$$|u(x, t) - v(x, t)| \leq \epsilon.$$

# Proof

- ▶ Let  $U(x, t) = u(x, t) - v(x, t)$ , then

$$U_t = u_t - v_t = ku_{xx} - kv_{xx} = kU_{xx}.$$

- ▶ For  $x = 0$ ,  $|U(0, t)| = |a_1(t) - a_2(t)| \leq \epsilon$  by assumption and thus

$$-\epsilon \leq U(0, t) \leq \epsilon \text{ for } 0 \leq t \leq T.$$

Likewise  $-\epsilon \leq U(L, t) \leq \epsilon$  for  $0 \leq t \leq T$ .

- ▶ For  $t = 0$ ,  $|U(x, 0)| = |f_1(x) - f_2(x)| \leq \epsilon$  by assumption and thus

$$-\epsilon \leq U(x, 0) \leq \epsilon \text{ for } 0 \leq x \leq L.$$

- ▶ Applying the Maximum and the Minimum Principles yields

$$-\epsilon \leq U(x, t) \leq \epsilon \text{ or}$$

$$|u(x, t) - v(x, t)| \leq \epsilon$$

for  $0 \leq x \leq L$  and  $0 \leq t \leq T$ .

# Uniqueness of Solutions

## Corollary

*Consider the initial boundary value problem*

$$\begin{aligned}u_t &= ku_{xx} + g(x, t), & 0 < x < L, & \quad t > 0 \\u(0, t) &= a(t) \quad \text{and} \quad u(L, t) = b(t), & 0 < x < L \\u(x, 0) &= f(x), & 0 \leq x \leq L\end{aligned}$$

*where the diffusion constant  $k > 0$ , the function  $g(x, t)$  is continuous, and the functions  $a(t)$ ,  $b(t)$ , and  $f(x)$  are  $C^2$  on their respective intervals. If there exists a  $C^2$  solution to this initial boundary value problem then it is unique.*

# Proof

- ▶ For the purposes of contradiction, suppose there are two solutions  $u(x, t)$  and  $v(x, t)$ .
- ▶ Define  $U(x, t) = u(x, t) - v(x, t)$ , then

$$\begin{aligned}U_t &= u_t - v_t \\&= ku_{xx} + g(x, t) - (kv_{xx} + g(x, t)) \\&= k(u_{xx} - v_{xx}) \\U_t &= kU_{xx} \text{ for } 0 < x < L \text{ and } t > 0.\end{aligned}$$

- ▶  $U(0, t) = u(0, t) - v(0, t) = a(t) - a(t) = 0$  and  $U(L, t) = 0$  and  $U(x, 0) = 0$
- ▶ By the Maximum and Minimum Principles  $U(x, t) = 0$  for all  $0 \leq x \leq L$  and  $t \geq 0$  which implies  $u(x, t) = v(x, t)$ .

# Homework

- ▶ Read Section 4.3
- ▶ Exercises: 21, 22, 23